

TECHNICAL SHEET

B1068	TTC BROTH BASE (TRICLOSAN TICARCILLIN CHLORATE BROTH)				
Formula					
Ingredients :		gms/lit.			
Casein enzymic hydrolysate		10.00			
Yeast extract		1.00			
Magnesium chloride, 6H ₂ O		60.00			
Sodium chloride		5.00			
Malachite green		0.01			
Trichlosan (Irgasan)		0.001			
Final pH (at 25°C) : 6.9 ± 0.2					
Directions :					
Suspend 44.11 grams (the equivalent weight of dehydrated medium per litre) in 988 ml distilled water. Heat to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Aseptically add 1 vial of each Ticarcillin Supplement (BF065) and Potassium Chlorate Supplement (BF066). Mix well before dispensing in sterile tubes or flasks as desired.					
Principle :					
Casein enzymic hydrolysate and yeast extract provides essential growth nutrients. Ticarcillin has inhibitory action on both gram – positive and gram – negative organisms. Irgasan inhibits gram – positive organisms. Potassium chlorate has disinfecting property.					
QC Tests – (I)Dehydrated Medium					
	Colour :		Light yellow to light blue		
	Appearance :		Homogeneous Free Flowing powder		
(II)Rehydrated medium					
	pH (post autoclaving/heating) :		6.9 ± 0.2		
	Colour (post autoclaving/heating) :		Peacock Green		
	Clarity (post autoclaving/heating) :		Clear		
(III)Q.C. Test Microbiological					
	Cultural characteristics observed with added Ticarcillin Supplement (BF065) and Potassium Chlorate Supplement (BF066) after an incubation at 25-30°C for 24-48 hours.				
	MICROORGANISM (ATCC)		GROWTH		
	Yersinia enterocolitica (27729)		Good-Luxuriant		
	Escherichia coli (25922)		inhibited		
	Staphylococcus aureus (25923)		inhibited		
Precautions :					
	1. For Laboratory Use.				
	2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :					
	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :					
	Recommended for selective enrichment and enumeration of Yersinia enterocolitica.				
Storage :					
	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :					
	100 gm. bottle /500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement
B1068		44.11 g/l	11.335 L	6.9 ± 0.2	Ticarcillin (BF065), Pot. Cholate (BF066)
					121°C/15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.