

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B1127	TRYPTONE BROTH (TRYPTONE WATER)				
Formula					
Ingredients :					
Casein enzymic hydrolysate	10.00				
Sodium chloride	5.00				
Final pH (at 25°C): 7.5 ± 0.2					
Directions :					
Suspend 15 grams in 1000 ml distilled water. Heat if necessary, to dissolve the medium completely. Dispense into tubes and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.					
Principle :					
Casein enzymic hydrolysate is a good substrate for indole production because of its high tryptophan content. Certain organisms breakdown the amino-acid tryptophan with the help of enzymes that mediate the production of indole by hydrolytic activity. The indole produced can be detected by either Kovac's or Ehrlich's reagent. Indole combines with the aldehyde present in the above reagent to give red colour in the alcohol layer. The alcohol layer extracts and concentrates the red colour complex.					
QC Tests - (I) Dehydrated Medium					
Colour :	Cream to yellow				
Appearance :	Homogeneous Free Flowing powder				
(II) Rehydrated medium					
pH (post autoclaving/heating) :	7.5 ± 0.2				
Colour (post autoclaving/heating) :	yellow				
Clarity (post autoclaving/heating) :	Clear				
(III) Q.C. Test Microbiological					
Cultural characteristics observed after an incubation at 35-37°C for 18-24 hours. Add 0.2 to 0.3ml of Kovac's Reagent (BA068) to each tube after incubation.					
MICROORGANISM (ATCC)	GROWTH	INDOLE REACTION			
Escherichia coli (25922)	Luxuriant	Positive reaction, red ring at the interface of the medium			
Enterobacter aerogenes (13048)	Luxuriant	Negative reaction, no colour development / cloudy ring			
Klebsiella pneumoniae (13383)	Luxuriant	Negative reaction, no colour development / cloudy ring			
Precautions :					
1. For Laboratory Use.					
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.					
Limitations :					
1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.					
Use :					
For the detection of indole production by coliforms from clinical and non-clinical samples.					
Storage :					
Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.					
Packing :					
500 gm. bottle					
Product profile:					
Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization	
B1127	15g/L	33.33L	7.5 ± 0.2	NIL	121°C /15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.