

TECHNICAL SHEET

B1271	TRYPTONE AGAR					
Formula						
Ingredients :			gms/lit.			
Casein enzymic hydrolysate			10.00			
Sodium chloride			8.00			
Agar			15.00			
Final pH (at 25°C) : Self						
Directions :						
Suspend 33.0 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour into sterile Petri plates.						
Principle :						
Casein enzymic hydrolysate provides essential growth nutrients to support the growth of organisms. Sodium chloride buffers the medium. This is also an excellent medium for the maintenance for both aerobic and anaerobic cultures. Viability in this medium is greater than in any other broth medium or slant culture						
QC Tests - (I) Dehydrated Medium						
Colour :			Cream to yellow			
Appearance :			Homogeneous Free Flowing powder			
(II) Rehydrated medium						
pH (post autoclaving/heating) :			Self			
Colour (post autoclaving/heating) :			Yellow			
Clarity (post autoclaving/heating) :			Clear to slightly opalescent			
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 24 - 48 hrs at 35-37°C.						
MICROORGANISM (ATCC)			GROWTH			
Enterobacter aerogenes (13048)			Luxuriant			
Escherichia coli (25922)			Luxuriant			
Salmonella enteritidis (13076)			Luxuriant			
Staphylococcus aureus (25923)			Luxuriant			
Pseudomonas aeruginosa (27853)			Luxuriant			
Precautions :		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		It is used as a general purpose medium for growth of non-fastidious microorganisms.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1271		33.0 g/l	15.15L	Self	Nil	121°C /15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.