

TECHNICAL SHEET

B1284	6 MFA MEDIUM				
Formula					
Ingredients :			gms/lit.		
Yeast extract	10.00				
Potassium dihydrogen phosphate	1.00				
Potassium chloride	0.50				
Sodium nitrate	3.00				
Magnesium sulphate	0.50				
Ferrous sulphate	0.01				
Zinc sulphate	0.001				
Sodium potassium tartrate	0.01				
Dextrose	40.00				
Agar	15.00				
Final pH (at 25°C) : Self					
Directions :					
Suspend 70.00 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Mix well and pour in sterile Petri plates.					
Principle :					
Yeast extract provides a source of trace elements, vitamins and amino acids. Dextrose is the source of carbohydrate. Monopotassium dihydrogen phosphate and sodium potassium tartarate buffers the media. Magnesium sulphate, ferrous sulphate and zinc sulphate are sources of divalent cations and other necessary ions. Agar is the solidifying agent.					
QC Tests - (I) Dehydrated Medium					
	Colour :		Light yellow to yellow		
	Appearance :		Homogeneous Free Flowing powder		
(II) Rehydrated medium					
	pH (post autoclaving/heating) :		Self		
	Colour (post autoclaving/heating) :		yellow		
	Clarity (post autoclaving/heating) :		clear to slightly opalescent gel		
(III) Q.C. Test Microbiological					
	Cultural characteristics observed after an incubation at 25-30°C for 24-48 hours.				
	MICROORGANISM (ATCC)		GROWTH		
	Aspergillus niger (16404)		good-luxuriant		
	Aspergillus ochraceous		good-luxuriant		
	Candida albicans (10231)		good-luxuriant		
	Saccharomyces cerevisiae (9763)		good-luxuriant		
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :	It is used for cultivation of <i>Aspergillus ochraceous</i>				
Storage :	Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1284	70.0 g/l	7.142 L	Self	NIL	121°C/ 15 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications. The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.