

**TECHNICAL SHEET**

<b>B1373</b>	<b>Mac Conkey Agar w/o sodium chloride</b>
<b>Formula</b>	
<b>Ingredients :</b>	<b>gms/lit.</b>
Peptic digest of animal tissue	20.00
Lactose	10.00
Bile salt	5.00
Neutral red	0.05
Agar	13.50
Final pH (at 25°C) : 7.4 ± 0.2	
<b>Directions :</b>	
Suspend 48.55 g of medium in 1000 ml. distilled water. Heat to boiling with gentle swirling to dissolve the agar completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Avoid overheating. Cool to 45 – 50°C and pour into sterile petri plates. The surface of the medium should be dry when inoculated.	
<b>Principle:</b> Peptic digest of animal tissue is sources of nitrogen and other nutrients. Lactose is a fermentable carbohydrate. When lactose is fermented, a local pH drop around the colony causes a colour change in the pH indicator (neutral red) Bile salt is selective agents that inhibit growth of gram –positive organisms. Agar is a solidifying agent.	

<b>QC Tests – (I)Dehydrated Medium</b>					
	Colour :	Pinkish beige			
	Appearance :	Homogeneous Free Flowing powder			
<b>(II)Rehydrated medium</b>					
	pH (post autoclaving/heating) :	7.4 ± 0.2			
	Colour (post autoclaving/heating) :	Orange red			
	Clarity (post autoclaving/heating) :	slightly opalescent			
<b>(III) Q.C. Test Microbiological</b>					
	Cultural characteristics observed after 18 – 24 hours at 35- 37°C.				
	MICROORGANISM (ATCC )	GROWTH	COLOUR OF COLONY		
	Enterobacter aerogenes (13048)	good-luxuriant	Pink to red		
	Escherichia coli (25922)	good-luxuriant	Pink to red with bile precipitate		
	Proteus vulgaris (13315 )	good-luxuriant	Colourless		
	Salmonella enteritidis (13076 )	good-luxuriant	Colourless		
	Salmonella typhi ( 6539 )	good-luxuriant	Colourless		
	Salmonella paratyphi A	good-luxuriant	Colourless		
	Salmonella paratyphi B	good-luxuriant	Colourless		
	Shigella flexneri (12022)	good-luxuriant	Colourless		
	Enterococcus faecalis (29212)	Fair to good	Pale pink to red		
	Staphylococcus aureus (25923)	Fair to good	Pale pink to red		
<b>Precautions :</b>		1. For Laboratory Use.			
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.			

Refer disclaimer overleaf

<b>Limitations :</b>	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
	2. Although MacConkey media are selective primarily for gram – negative enteric bacilli, biochemical and, if indicated, serological testing using pure cultures are recommended for complete identification. Consult appropriate references for further information.				
	3. Incubation of MacConkey Agar plates under increased CO <sub>2</sub> has been reported to reduce the growth and recovery of a number of strains of gram – negative bacilli.				
<b>Use :</b>	For cultivation and differentiation of enteric bacteria, restricting swarming of Proteus species from specimens like urine which may contain large number of Proteus species as well as potentially pathogenic gram-positive organisms.				
<b>Storage :</b>	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
<b>Packing :</b>	500 gm. bottle				
<b>Product profile:</b>	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B1373</b>	48.50g/L	10.309L	7.4± 0.2	NIL	121°C /15 min.

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.