

TECHNICAL SHEET

B1445	TRYPTIC SOYA AGAR				
Formula					
Ingredients :		gms/lit.			
Pancreatic digest of casein	17.00				
Papaic digest of soyabean meal	3.00				
Sodium chloride	5.00				
Dextrose (Glucose)	2.50				
Dipotassium hydrogen phosphate	2.50				
Agar	15.000				
Final pH (at 25°C): 7.3 ± 0.2					
Directions :					
Suspend 45 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates.					
Principle :					
Pancreatic digest of casein and papaic digest of soyabean meal provides nitrogen, vitamins and minerals. Glucose serves as the carbohydrate source. Phosphate buffers the media. Sodium chloride maintains the osmotic balance. Agar is the solidifying agent.					
QC Tests – (I)Dehydrated Medium					
	Colour :	Cream to yellow			
	Appearance :	Homogeneous Free Flowing powder			
(II)Rehydrated medium					
	pH (post autoclaving/heating) :	7.3 ± 0.2			
	Colour (post autoclaving/heating) :	Light yellow			
	Clarity (post autoclaving/heating) :	clear to slightly opalescent gel			
(III)Q.C. Test Microbiological					
	Cultural characteristics was observed after an incubation at 30-35°C 18-24 hours				
	MICROORGANISM (ATCC)	GROWTH			
	Salmonella Typhi (6539)	Luxuriant			
	Salmonella Typhimurium (14028)	Luxuriant			
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :	It is used for general purpose medium used for a cultivation and maintenance of Salmonella typhimurium.				
Storage :	Dehydrated medium-below 30°C Prepared medium- Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B1445	45.0 g/l	11.111 L	7.3 ± 0.2	Nil	121°C /15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.