

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B1805	ANAEROBIC BROTH	
Formula		
Ingredients :	gms/lit.	
Tryptone	20.00	
Dextrose(Glucose)	10.00	
Sodium chloride	5.00	
Sodium thioglycollate	2.00	
Sodium formaldehyde Sulfoxylate	1.00	
Methylene blue	0.002	
Final pH (at 25°C) : 7.2 ± 0.2		
Directions :		
Suspend 38.0 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.		
Principle :		
The medium contains sodium thioglycollate and Sodium formaldehyde Sulfoxylate that provide adequate anaerobiosis which is indicated by methylene blue present in the medium which yields blue colour to medium in presence of oxygen. Tryptone and dextrose provide essential nutrients while sodium chloride maintains osmotic equilibrium.		
QC Tests – (I)Dehydrated Medium		
Colour :	Cream to yellow	
Appearance :	Homogeneous Free Flowing powder	
(II)Rehydrated medium		
pH (post autoclaving/heating) :	7.2 ± 0.2	
Colour (post autoclaving/heating) :	Light green	
Clarity (post autoclaving/heating) :	Clear	
(III)Q.C. Test Microbiological		
Cultural characteristics observed after 48-72 hrs. at 35-37°C when incubated anaerobically.		
MICROORGANISM (ATCC)	GROWTH	
Clostridium butyricum (13732)	Good-luxuriant	
Clostridium perfringens (12924)	Good-luxuriant	
Clostridium sporogenes (11437)	Good-luxuriant	
Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.	
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Clinical specimens must be obtained properly and transported to the laboratory in a suitable anaerobic transport container. 3. The microbiologist must be able to verify quality control of the medium and determine whether the environment is anaerobic. 4. The microbiologist must perform aerotolerance testing on each isolate recovered to ensure that the organism is an anaerobe. 5. Methylene blue is toxic to some anaerobic bacteria.	
Use :	For the cultivation of anaerobic bacteria, especially Clostridium species and other anaerobic organisms from clinical and non-clinical samples.	
Storage :	Dehydrated medium-below 30°C Prepared medium- Between 20 to 30°C.	
Packing :	500 gm. bottle	

Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B104	38.00 g/l	13.15 L	7.2± 0.2	Nil	121°C /15 min.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.