

**TECHNICAL SHEET**

<b>B220</b>	<b>LAURYL SULPHATE BROTH (LAURYL TRYPTOSE BROTH)</b>				
<b>Formula</b>					
<b>Ingredients:</b>		<b>gms/lit.</b>			
Tryptose		20.00			
Lactose		5.00			
Sodium chloride		5.00			
Dipotassium phosphate		2.75			
Monopotassium phosphate		2.75			
Sodium lauryl sulphate		0.10			
Final pH (at 25°C): 6.8 + 0.2					
<b>Directions:</b>					
Suspend 35.6 grams in 1000 ml distilled water. Heat, if necessary, to dissolve the medium completely. Distribute into tubes containing inverted Durham’s tubes. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. For inoculum of 1 ml or less, use single strength medium. For inocula of 10 ml or more, double strength or proportionate medium should be prepared.					
<b>Principle:</b>					
Tryptose provides essential growth substances, such as nitrogen and carbon compounds, sulphate and trace ingredients. The potassium phosphates provide buffering system, while sodium chloride maintains osmotic equilibrium. Sodium lauryl sulphate inhibits organisms other than coliforms. Lactose is the fermentable carbohydrate for coliforms.					
<b>QC Tests – (I)Dehydrated Medium</b>					
	Colour :		Cream to light yellow		
	Appearance:		Homogeneous Free Flowing powder		
<b>(II)Rehydrated medium</b>					
	pH (post autoclaving/heating):		6.8 ± 0.2		
	Colour (post autoclaving/heating):		Light amber to light yellow		
	Clarity (post autoclaving/heating):		Clear		
<b>(III) Q.C. Test Microbiological</b>					
	Cultural characteristics observed after18- 24 hours at 35-37°C.				
	MICROORGANISM (ATCC)	GROWTH	GAS PRODUCTION	INDOLE (44°C)	
	Enterobacter aerogenes (13048)	Luxuriant	+	negative reaction, no colour development / cloudy ring	
	Escherichia coli (25922)	Luxuriant	+	positive reaction, red ring at the interface of the medium	
	Salmonella typhimurium (14028)	Luxuriant	-	negative reaction, no colour development / cloudy ring	
	Staphylococcus aureus (25923)	Inhibited	-	-	
	Enterococcus faecalis (29212)	Inhibited	-	-	
<b>Precautions :</b>		1. For Laboratory Use.			
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.			
<b>Limitations :</b>		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.			
<b>Use:</b>		For detection and enumeration of coliform bacteria in water, wastewater, dairy products and other foods.			
<b>Storage:</b>		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.			
<b>Packing:</b>		500 gm. bottle			
<b>Product profile:</b>		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement Sterilization
<b>B220</b>		35.6 g/l	14.04 L	6.8 ± 0.2	NIL 121 <sup>o</sup> C/15 MIN

Refer disclaimer Overleaf

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.