

## TECHNICAL SHEET

B233		MIO MEDIUM (MOTILITY INDOLE ORNITHINE MEDIUM)				
Formula						
Ingredients :		gms/lit.				
Casein enzymic hydrolysate		10.00				
Peptic digest of animal tissue		10.00				
Yeast extract		3.00				
L-Ornithine hydrochloride		5.00				
Dextrose		1.00				
Bromo cresol purple		0.02				
Agar		2.00				
Final pH (at 25°C) : 6.5 ± 0.2						
Directions :						
Suspend 31.02 grams in 1000 ml distilled water. Heat to boiling to dissolve the medium completely. Dispense in test tubes in 5 ml amounts. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubes in an upright position.						
Principle :						
MIO Medium contains Peptic digest of animal tissue and Casein enzyme hydrolysate which provide carbon and nitrogen. Yeast Extract provides vitamins and cofactors required for growth as well as additional sources of nitrogen and carbon. Dextrose is an energy source. Agar is added to demonstrate motility. The pH indicator, Bromo Cresol Purple, facilitates detection of decarboxylase activity.						
QC Tests – (I)Dehydrated Medium						
Colour :		Light yellow to pale green				
Appearance :		Homogeneous Free Flowing powder				
(II)Rehydrated medium						
pH (post autoclaving/heating) :		6.5 ± 0.2				
Colour (post autoclaving/heating) :		Purple				
Clarity (post autoclaving/heating) :		Clear to slightly opalescent				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 40 – 48 hours at 35 - 37°C.						
MICROORGANISM (ATCC )		GROWTH	MOTILITY	INDOLE	ORNITHINE DECARBOXYLASE	
Enterobacter aerogenes (13048)		Luxuriant	+	-	+	
Escherichia coli (25922)		Luxuriant	+	+	+	
Klebsiella pneumoniae (13883 )		Luxuriant	-	-	-	
Proteus mirabilis (25933)*		Luxuriant	+	-	+	
Key : + = positive reaction						
- = negative reaction						
* = motility of Proteus mirabilis is temperature dependent. It is more Pronounced at 20°C and almost absent at 35°C.						
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For the identification of Enterobacteriaceae on the basis of motility, indole production and ornithine decarboxylase activity.				
Storage :		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B233		31.02 g/l	16.118 L	6.5 ± 0.2	NIL	121 <sup>o</sup> C / 15 minutes

Refer disclaimer Overleaf

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.