

**TECHNICAL SHEET**

<b>B263</b>	<b>MUELLER HINTON AGAR</b>	
<b>Formula</b>		
<b>Ingredients :</b>	<b>gms/lit.</b>	
Meat, infusion from	300.00	
Casein acid hydrolysate	17.50	
Starch	1.50	
Agar	17.00	
# Equivalent to beef, infusion form		
Final pH (at 25°C) :7.3± 0.1		
<b>Directions :</b>		
Suspend 38.0 grams in 1000 ml purified/ distilled water. Heat to boiling to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool to 45-50°C. Mix well and pour into sterile Petri plates		
Note: The performance of this batch has been tested and standardized as per the current CLSI (formerly, NCCLS) document M6-protocols for Evaluating Dehydrated Mueller Hinton Agar.		
<b>Principle :</b>		
Meat, infusion from and casein acid hydrolysate provide nitrogenous compounds, carbon, sulphur and other essential nutrients. Starch is added to absorb any toxic substances present in the medium. Different factors influence the disc diffusion susceptibility tests as, inoculum concentration, agar depth, disc potency, medium pH and beta – lactamase production by test organisms. Agar is the solidifying agent.		
<b>QC Tests – (I)Dehydrated Medium</b>		
	Colour :	Cream to yellow
	Appearance :	Homogeneous Free Flowing powder
<b>(II)Rehydrated medium</b>		
	pH (post autoclaving/heating) :	7.3 ± 0.1
	Colour (post autoclaving/heating) :	Light amber
	Clarity (post autoclaving/heating) :	Slightly opalescent
<b>(III) Q.C. Test Microbiological</b>		
	Cultural characteristics observed after 18- 24 hours at 36-37°C.	
	MICROORGANISM (ATCC )	GROWTH
	Escherichia coli (25922)	Luxuriant
	Pseudomonas aeruginosa (27853)	Luxuriant
	Staphylococcus aureus (25923)	Luxuriant
	Enterococcus faecalis(29212)	Luxuriant
	Escherichia coli (35218)	Luxuriant
	Staphylococcus aureus subsp. aureus (43300)	Luxuriant
<b>Precautions :</b>	1. For Laboratory Use.	
	2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.	

Refer disclaimer Overleaf

**TECHNICAL SHEET**

<b>Limitations :</b>	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
	2. Numerous factors can affect results ; inoculum size, rate of growth, medium formulation of pH, length of incubation and incubation environment, disk content and drug diffusion rate, and measurement of endpoints. Therefore, strict adherence to protocol is required to ensure reliable results.				
	3. Disk diffusion susceptibility testing is limited to rapidly growing organisms. Drug inactivation may result from the prolonged incubation times required by slow growers.				
	4. Media containing excessive amounts of thymidine or thymine can reverse the inhibitory effects of sulfonamides and trimethoprim, causing zones of growth inhibition to be smaller or less distinct.				
	5. Variation in the concentration of divalent cations, primarily calcium and magnesium, affects results of aminoglycoside, tetracycline, and colistin tests with <i>P. aeruginosa</i> isolates. A cation content that is too high reduces zones sizes, whereas a cation content that is too low has the opposite effect.				
	6. When Mueller Hinton Medium is supplemented with blood, the Zone of inhibition for oxacillin and methicillin may be 2 to 3 mm smaller than those obtained with unsupplemented agar. Conversely, sheep blood may markedly increase the zone diameters of some cephalosporins when they are tested against enterococci. Sheep blood may cause indistinct zones or a film of growth within the zones of inhibition around sulfonamide and trimethoprim disks.				
	7. Mueller Hinton Medium deeper than 4 mm may cause false – resistant results, and agar less than 4 mm deep may be associated with a false – susceptibility report.				
	8. A pH outside the range of $7.3 \pm 0.1$ may adversely affect susceptibility test results. If the pH is too low, aminoglycosides and macrolides will appear to lose potency; others may appear to have excessive activity. The opposite effects are possible if the pH is too high.				
	9. When Mueller Hinton Medium is inoculated, no droplets of moisture should be visible on the surface or on the petri dish cover.				
	10. Mueller Hinton Medium should be inoculated within 15 minutes after the inoculum suspension has been adjusted.				
	11. The zone of inhibition diameters of some drugs, such as the aminoglycosides, macrolides, and tetracyclines, are significantly altered by CO <sub>2</sub> Plates should not be incubated in increased CO <sub>2</sub> .				
	12. This medium is recommended for susceptibility testing of pure cultures only.				
<b>Use:</b>	Recommended for determination of susceptibility of microorganisms to antimicrobial agents isolated from clinical samples.				
<b>Storage :</b>	Dehydrated medium -below 30°C Prepared medium – Between 2 to 8°C.				
<b>Packing :</b>	500 gm. bottle				
<b>Product profile:</b>	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
<b>B263</b>	38g/l	13.157L	7.3 ± 0.1	NIL	121°C / 15 minutes

**Disclaimer:**

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.