

TECHNICAL SHEET

B350	VIOLET RED BILE AGAR				
Formula					
Ingredients :		gms/lit.			
Peptic digest of animal tissue		7.00			
Yeast extract		3.00			
Lactose		10.00			
Bile salt mixture		1.50			
Sodium chloride		5.00			
Neutral red		0.03			
Crystal violet		0.002			
Agar		15.00			
Final pH (at 25°C) : 7.4 ± 0.2					
Directions :					
Suspend 41.53 grams in 1000 ml distilled water. Heat with stirring to boiling to dissolve the medium completely. DO NOT AUTOCLAVE. Cool to 45°C and pour into sterile petri plates containing the inoculum. If desired, the medium can be sterilized by autoclaving at 15 lbs pressure (121°C) for 15 minutes.					
Principle :					
Violet Red Bile Agar (VRBA) contains peptic digest of animal tissue to provide carbon and nitrogen sources for general growth requirements, Yeast extract supplies B – complex vitamins which stimulate bacterial growth. Bile salts and Crystal Violet inhibit most gram– positive microorganisms. Lactose is the carbohydrate source and Neutral red is the pH indicator, Agar is the solidifying agent.					
QC Tests – (I)Dehydrated Medium					
	Colour :	Light yellow to pink			
	Appearance :	Homogeneous Free Flowing powder			
(II)Rehydrated medium					
	pH (post autoclaving/heating) :	7.4 ± 0.2			
	Colour (post autoclaving/heating) :	Reddish purple			
	Clarity (post autoclaving/heating) :	Clear to slightly opalescent			
(III)Q.C. Test Microbiological					
	Cultural characteristics observed after 18 - 24 hrs. at 35-37°C.				
	MICROORGANISM (ATCC)	GROWTH	COLOUR OF COLONY		
	Escherichia coli (25922)	Luxuriant	Pinkish red with bile precipitate		
	Enterobacter aerogenes (13048)	Luxuriant	Pink to pinkish red		
	Salmonella enteritidis (13076)	Luxuriant	Colourless to orangish yellow		
	Staphylococcus aureus (25923)	Inhibited	--		
Precautions:	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Violet Red Bile Agar may not be completely inhibitory to gram – positive organisms. Perform Gram stain and biochemical tests as necessary to identify isolates. 3. The medium will grow gram- negative bacilli other than members of the Enterobacteriaceae. Perform biochemical tests to identify isolates to genus and species. 4. Boiling the medium for longer than 2 minutes can decrease the ability to support growth. 5. Plates of Violet Red Bile Agar should not be incubated longer than 24 hours because microorganisms that are only partially inhibited may grow after extended incubation. 6. For optimum performance, prepare and use the medium within 24 hours.				
Use:	For selective isolation, detection and enumeration of coli-aerogenes bacteria in water, milk dairy and other food products.				
Storage:	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B350	41.53 g/l	12.03L	7.4 ± 0.2	NIL	DO NOT AUTOCLAVE

Refer disclaimer overleaf

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.