

TECHNICAL SHEET

B361	XYLOSE LYSINE DEOXYCHOLATE AGAR (XLD AGAR)		
Formula			
Ingredients :	gms/lit.		
Yeast extract	3.00		
L-Lysine	5.00		
Lactose	7.50		
Sucrose	7.50		
Xylose	3.50		
Sodium chloride	5.00		
Sodium deoxycholate	2.50		
Sodium thiosulphate	6.80		
Ferric ammonium citrate	0.80		
Phenol red	0.08		
Agar	15.00		
Final pH (at 25°C) : 7.4 ± 0.2			
Directions :			
Suspend 56.68 grams in 1000 ml distilled water. Heat with frequent agitation until the medium boils. DO NOT AUTOCLAVE OR OVERHEAT. Transfer immediately to a water bath at 50°C. After cooling, pour into sterile Petri plates. It is advisable not to prepare large volumes that will require prolonged heating, thereby producing precipitate.			
Note: Slight precipitation in the medium may occur, which is inheritant property of the medium, and does not affect the performance of the medium.			
Principle :			
Yeast extract provides sources of nitrogen and carbon, as well as vitamins and cofactors required for growth. Xylose, lactose, and sucrose (Saccharose) are fermentable carbohydrates. Xylose is fermented by most enteric organisms except Shigella and Providencia. Lysine is added to differentiate Salmonella. As xylose is exhausted, Salmonella organisms decarboxylate lysine causing reversion to alkaline conditions. Alkaline reversion by other lysine – positive organisms is prevented by excess acid production from fermentation of lactose and sucrose.			
Sodium Thiosulfate and Ferric Ammonium citrate allow visualization of hydrogen sulfide production under alkaline conditions. Acidic conditions inhibit the reaction. Phenol red is an indicator. Sodium chloride maintains osmotic balance in the medium. Agar is a solidifying agent.			
Sodium Deoxycholate in XLD agar inhibits growth of gram – positive organisms.			
QC Tests – (I)Dehydrated Medium			
Colour :	Light yellow to pink		
Appearance :	Homogeneous Free Flowing powder		
(II)Rehydrated medium			
pH (post autoclaving/heating) :	7.4 ± 0.2		
Colour (post autoclaving/heating) :	Red		
Clarity (post autoclaving/heating) :	Clear to very slightly opalescent		
(III)Q.C. Test Microbiological			
Cultural characteristics observed after 18 - 72 hrs. at 35 - 37°C.			
MICROORGANISM (ATCC)	GROWTH	COLOUR OF COLONY	INCUBATION PERIOD
Proteus vulgaris (13315)	Good –luxuriant	grey with black centres	18 -72 hrs
Salmonella enteritidis (13076)	Good –luxuriant	Red with black centers	18 -72 hrs
Salmonella paratyphi A	Good –luxuriant	Red	18 -72 hrs
Salmonella paratyphi B	Good –luxuriant	Red with black centers	18 -72 hrs
Salmonella typhi (6539)	Good –luxuriant	Red with black centers	18 -72 hrs
Salmonella typhimurium (14028)	Luxuriant	Red with black centers	18 -72 hrs
Salmonella Abony (NCTC6017)	good-luxuriant	Red with black centers	18 -72 hrs
Shigella sonnei (25931)	fair-good	Red	18 -72 hrs
Shigella dysenteriae (13313)	Good –luxuriant	Red	18 -72 hrs

Refer disclaimer Overleaf

TECHNICAL SHEET

Shigella flexneri (12022)	fair-good	Red	18 -72 hrs
Enterobacter aerogenes (13048)	Fair	Yellow	18 -72 hrs
Escherichia coli (25922)	Fair	Yellow	18 -72 hrs
Escherichia coli (8739)	Fair	Yellow	18 -72 hrs
Escherichia coli (NCTC9002)	Fair	Yellow	18 -72 hrs
Staphylococcus aureus (25923)	inhibited	-	>=72 hrs
Staphylococcus aureus (8539)	inhibited	-	>=72 hrs
Enterococcus faecalis (29212)	inhibited	-	>=72 hrs

Precautions :	1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium. 2. Non-enterics like Pseudomonas and Providencia may exhibit red colonies. 3. S. paratyphi A, S. choleraesuis, S. pullorum and S. gallinarum may form red colonies without black centers, thus resembling Shigella species. 4. Some Proteus strains will give black – centered colonies on XLD Agar. 5. Slight precipitation in the medium may occur, which is inheritant property of the medium, and does not affect the performance of the medium.				
Use :	For selective isolation and enumeration of Salmonella typhi and other Salmonella species.				
Storage :	Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :	500 gm. bottle				
Product profile:	Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B361	56.68 g/l	8.82 L	7.4 ± 0.2	Nil	Do not autoclave/ over heat. Boil medium to dissolve w/frequent agitation

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.