

TECHNICAL SHEET

B723	SKIM MILK					
Formula						
Ingredients :			gms/lit.			
Skim milk powder			100.00			
Final pH (at 25°C) :			7.0 ± 0.2			
Directions :						
Suspend 100 gms in a little amounts of distilled water to make a smooth paste. Gradually add more distilled water to make a final volume of 1000 ml. Dispense & sterilize by autoclaving at 15 lbs pressure (121°C) for 5 minutes.						
Principle :						
Skim milk is a source of lactose and casein. Hydrolysis of casein is detected by visible formation of a clot. Skim milk serves as a good source of casein.						
QC Tests – (I)Dehydrated Medium						
	Colour :		Cream to beige			
	Appearance :		Homogeneous Free Flowing powder			
(II)Rehydrated medium						
	pH (post autoclaving/heating) :		7.0 ± 0.2			
	Colour (post autoclaving/heating) :		White to off white			
	Clarity (post autoclaving/heating) :		Opaque			
(III)Q.C. Test Microbiological						
	Cultural characteristics observed after 18 – 24 hrs. at 35- 37°C.					
	MICROORGANISM (ATCC)	GROWTH	PROTEOLYTIC ACTIVITY			
	Bacillus subtilis (6633)	Luxuriant	+			
	Escherichia coli (25922)	Luxuriant	-			
	Pseudomonas aeruginosa (27853)	Luxuriant	+			
	Proteus mirabilis (25933)	Luxuriant	+			
	Enterococcus faecalis (29212)	Luxuriant	-			
	Serratia marcescens (8100)	Luxuriant	+			
Precautions :		1. For Laboratory Use.				
		2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
		2. Skim Milk supports growth of many microorganisms. Perform microscopic examination and other biochemical tests to identify isolates to the genus and species level, if necessary.				
Use :		For cultivation of dairy organisms and differentiation of Clostridium species.				
Storage :		Dehydrated medium- below 30°C Prepared medium– Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B723		100g/l	5L	7.0 ± 0.2	NIL	121°C / 5 minutes

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents