

B821	YEAST EXTRACT AGAR					
Formula						
Ingredients :		gms/lit.				
Peptic digest of animal tissue		5.00				
Yeast extract		3.00				
Agar		15.00				
Final pH (at 25°C) : 7.2 ± 0.2						
Directions :						
Suspend 23 gms. in 1000 ml. distilled water. Boil to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.						
Principle :						
Yeast extract and peptic digest of animal tissue provide nitrogenous compounds, vitamin B complex and other growth nutrients. From the water sample, make a decimal dilution bank with Ringer Solution (B381) and take aliquots to 2 parallel series of plates. Pour the molten, cooled (45°C) Yeast Extract Agar and homogenize with sample. Incubate one of the series of plates at 35°C for 24 hours and the other series of plates at 20-22°C for 3 days. Separate counts are made of the organisms forming visible colonies after 24 hours at 35°C and the organisms forming colonies after 3 days at 20-22°C. Select the plates containing 30-300 colonies.						
QC Tests - (I) Dehydrated Medium						
Colour :		Cream to light yellow				
Appearance :		Homogeneous Free Flowing powder				
(II) Rehydrated medium						
pH (post autoclaving/heating) :		7.2 ± 0.2				
Colour (post autoclaving/heating) :		Light yellow to yellow				
Clarity (post autoclaving/heating) :		Clear to very slightly opalescent				
(III) Q.C. Test Microbiological						
Cultural characteristics observed after 18 - 24 hrs at 35 - 37°C.						
MICROORGANISM (ATCC)		GROWTH				
Escherichia coli (25922)		Luxuriant				
Staphylococcus aureus (25923)		Luxuriant				
Enterobacter aerogenes (13048)		Luxuriant				
Pseudomonas aeruginosa (27853)		Luxuriant				
Precautions :		1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.				
Limitations :		1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.				
Use :		For plate count of microorganisms in water.				
Storage :		Dehydrated medium- below 30°C Prepared medium- Between 2 to 8°C.				
Packing :		500 gm. bottle				
Product profile:		Reconstitution	Quantity on Preparation (500g)	pH (25°C)	Supplement	Sterilization
B821	23 g/l	21.73L	7.2 ± 0.2	Nil	121°C/15 min.	

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARK LABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.