

BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

B910	ACETAMIDE AGAR (TWIN PACK)				
Formula					
Ingredient:				gms/lit.	
Part A:					
Acetamide					10.00
Part B:					
Sodium chloride					5.00
Dipotassium hydrogen phosphate					1.39
Potassium dihydrogen phosphate					0.73
Magnesium sulphate					0.50
Phenol red					0.012
Agar					15.00
Final pH (at 25°C):					7.0 ± 0.2
Directions:					
Suspend 22.63 grams of part B in 1000 ml distilled water. Add 10.0 grams of Part A. Heat to boiling to dissolve the medium completely. Dispense in tubes or as desired. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. Cool the tubes in a slanted position.					
Principle :					
The medium contains inorganic salts and acetamide a sole carbon and nitrogen source. Sodium chloride maintains the osmotic equilibrium. Phenol red is the pH indicator. Organisms growing in the medium metabolize acetamide by the process of deamination (acylamidase activity). This ability is shown by <i>Pseudomonas aeruginosa</i> , <i>Pseudomonas acidovorans</i> Group III (<i>Achromobacter xylosoxidans</i>) and <i>Alcaligenes odorans</i> . Deamination of acetamide produces ammonia which increases the pH of the media causing a corresponding colour change from yellow - orange to purplish red. Some strains deamidate acetamide slowly and may require upto 7 days.					
QC Tests – (I)Dehydrated Medium					
Colour :		Part A) Colourless		Part B) Light yellow to brick red	
Appearance :		Part A) Deliquescent crystals Part B) Homogeneous Free Flowing powder			
(II)Rehydrated medium					
pH (post autoclaving/heating) :		7.0 ± 0.2			
Colour (post autoclaving/heating) :		Orange			
Clarity (post autoclaving/heating) :		Clear to slightly opalescent			
(III)Q.C. Test Microbiological					
Cultural characteristics observed after an incubation at 35-37°C for 4-7 days.					
MICROORGANISM (ATCC)		GROWTH		DEAMINATION	
Pseudomonas aeruginosa 27853)		Good -luxuriant		Positive reaction, purplish red colour within 7 days	
Pseudomonas maltophilia (13637)		Good -luxuriant		Negative reaction ,no purplish red colour within 7 days	
Precautions :					
1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.					
Limitations :					
1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.					
Use:					
For confirmation of <i>Pseudomonas aeruginosa</i> in water samples.					
Storage:					
Dehydrated medium-below 30°C Prepared medium- Between 2 to 8°C.					
Packing :					
500 gm. bottle					
Product profile:		Reconstitution		Quantity on Preparation (500g)	pH (25°C)
		Supplement		Sterilization	
B910		22.63 g/l part A 10.00 g/l part B		15.32 L 28.36 L	7.0 ± 0.2
		None		121°C/15 min.	

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.