

**BF059**

**V.C.N. SUPPLEMENT**

**FORMULA**

**Ingredients :**

(Per vial, sufficient for 500 ml medium)

Vancomycin	1.500mg
Colistin methane sulphonate	3.750mg
Nystatin	6250Units

**Directions :** Rehydrate the contents of 1 vial aseptically with 2 ml sterile distilled water. Mix well and aseptically add it to Thayer Martin Medium Base (B333) for 440 ml of medium aseptically add 50 ml sterile lysed blood and one vial of V.C.N. Supplement BF059 along with one vial of Vitamino Growth Supplement (BF037) 2% Haemoglobin Solution (250ml) (BF036) can be used instead of sterile lysed blood in 250 ml of medium. In GC Agar Base (B1049) for 250 ml of medium aseptically add 250 ml of 2% Haemoglobin Solution (BF036) and GC Supplement w/Antibiotics (BF058) one vial of Vitamino Growth Supplement (BF037) or Yeast Autolysate Supplement (BF038) If desired V.C.N. Supplement (BF059) can be used along with GC Supplement w/Antibiotics (BF058) for additional selectivity. In Transgrow Medium Base (B078) for 440 ml of medium aseptically add 50 ml of sterile 2% Haemoglobin Solution (BF036) and one vial of V.C.N. Supplement (BF059) along with one vial of Vitamino Growth Supplement (BF037)

**Precautions :**

1. For Laboratory Use.
2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.

**Use :** An antibiotic supplement, recommended for the selective isolation of Neisseria meningitidis and Neisseria meningitidis

**Storage:** Between 2 to 8°C.

**Packing :** 5 vial