

<b>BF062</b>	<b>VANCO T SUPPLEMENT</b> (vancomycin-colistin-amphotericin b-trimethoprim)
<b>FORMULA</b>	
<b>Ingredients :</b>	
(per vial sufficient for 500 ml. of medium)	
Vancomycin	1.00 mg
Celestin sulphate	3.75 mg
Amphotericin B	0.50 mg
Trimethoprim lactate	1.50 mg
<b>Directions:</b>	
Rehydrate the contents of 1 vial aseptically with 10 ml sterile distilled water. Mix gently, avoiding frothing to dissolve. Aseptically add the contents to 490 ml sterile, molten GC agar base (B1049) containing 50 ml sterile defibrinated horse blood lysed with 0.5% (v/v) saponin and rehydrated contents of 1 vial of Yeast autolysate supplement (BF038). Mix well and pour into sterile Petri plates.	
<b>Precautions :</b>	
1. For Laboratory Use. 2. Follow proper, established laboratory procedures in handling and disposing of infectious materials.	
<b>Use :</b>	An antibiotic supplement recommended for the selective isolation of <i>Nesseria gonorrhoeae</i> and <i>Neisseria meningitidis</i> .
<b>Storage:</b>	Between 2 to 8°C.
<b>Packing :</b>	1 vial