

TECHNICAL SHEET

PP006	CLED Agar w/Bromo Thymol Blue Plate				
Formula					
Ingredients:		gms/lit.			
Peptic digest of animal tissue		4.00			
Casein enzymic hydrolysate		4.00			
Meat Extract B#		3.00			
Lactose		10.00			
L-cystine		0.128			
Bromo thymol blue		0.02			
Agar		15.00			
#- Equivalent to Beef extract					
Final pH (at 25°C): 7.3 ± 0.2					
Directions:					
Label the ready to use plate (PP006). Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.					
Principle:					
Beef Extract, Peptic digest of animal tissue and Casein enzyme hydrolysate provide the nitrogen, vitamins and amino acids in CLED Agar, L-Cystine is added as a growth supplement for cystine – dependent coliforms. Lactose is included as a carbon source. Organisms capable of fermenting lactose will lower the pH and change the color of the medium from green to yellow. Bromo Thymol Blue is used as a pH indicator. Shigella species may not grow on this medium.					
(I) QC Tests					
pH:		7.3 ± 0.2			
Color:		Green coloured medium.			
Appearance:		Sterile CLED Agar w/Bromo Thymol Blue in 90X15mm disposable plates.			
(II) Sterility test		Passes release criteria			
(III) Q.C. Test Microbiological					
Cultural characteristics observed after incubation at 35-37°C for 18-24 hours.					
	MICROORGANISM (ATCC)	INOCULUM (CFU)	GROWTH	RECOVERY	COLOR OF COLONY
	Enterococcus faecalis 29212	50-100	good-luxuriant	≥70%	slight yellowish or greenish
	Klebsiella pneumoniae 13883	50-100	good-luxuriant	≥70%	yellow to whitish blue
	Escherichia coli ATCC 25922	50-100	good-luxuriant	≥70%	yellow, opaque, centre slightly deeper yellow
	Proteus vulgaris 13315	50-100	good-luxuriant	≥70%	blue
	Salmonella Typhi 6539	50-100	good-luxuriant	≥70%	bluish
	Staphylococcus aureus 25923	50-100	good-luxuriant	≥70%	deep yellow

TECHNICAL SHEET

Precautions :	1. In Vitro diagnostic use only.
	2. Read the label before opening the container
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.
Use:	For isolation and differentiation of urinary pathogens on the basis of lactose fermentation.
Storage:	Store between 2-8°C. Use before expiry date on the label.
Packing:	20/50 disposable plates.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.