BIOMARK Laboratories-INDIA

www.biomarklabs.com

TECHNICAL SHEET

PP013 MacCo	onkey Sorbitol Agar Plate			
Formula				
Ingredients:	gms/lit.			
Tryptone	17.00			
Meat Peptone	3.00			
D-Sorbitol	10.00			
Bile salt No.3	1.50			
Sodium chloride	5.00			
Neutral red	0.03			
Crystal violet	0.001			
Agar	13.50			
Final pH (at 25°C): 7	.1 <u>+</u> 0.2			
Directions:				

Label the ready to use plate (PP013). Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.

Principle:

Tryptone and Meat Peptone are sources of nitrogen and other nutrients. D-Sorbitol is a fermentable carbohydrate. Crystal Violet and Bile Salts are selective agents that inhibit growth of gram –positive organisms. Agar is a solidifying agent. The growth of E. coli O157:H7 on MacConkey Agar with Sorbitolshows colorless colonies and most of the fecal flora ferment sorbitol and appear pink. MacConkey Agar with Sorbitol therefore permits ready recognition of E. coli O157: H7which ferments lactose but does not ferment sorbitol, henceproduces colorless colonies.

CC	noness colonies.				
(I)	(I) QC Tests				
	pH:	7.1 ± 0.2			
	Color:	Purplish red coloured me	edium		
	Appearance:	Sterile MacConkey Sort	oitol Agar in 90X15 mm		
		disposable plate.			
(II)Sterility test		Passes release criteria	1		
(I	II)Q.C. Test Microbiological				
	Cultural response wasobserved	after an incubation a	at 30-35°C for 18-72		
	hours.				
	MICROORGANISM (ATCC)	GROWTH	COLOR OFCOLONY		
	Escherichia coli 25922	luxuriant	Pale pink-red		
	Escherichia coli serotype011	luxuriant	Pale pink-red		
	and 055				
	Escherichia coli 0157:H7	luxuriant	Colorless		
	S. serotype typhi 6539	Fair to good	Colorless		
	Shigella flexneri 12022	luxuriant	Colorless		

Refer disclaimer Overleaf Page 01 of 02

Rev: January2025

BIOMARK Laboratories-INDIA www.biomarklabs.com

TECHNICAL SHEET

Precautions:	In Vitro diagnostic use only.		
	2. Read the label before opening the container		
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be		
	encountered that fail to grow or grow poorly on this medium.		
Use:	for isolation and identification of enteropathogenic Escherichia coli strains		
	associated with infant diarrhoea.		
Storage:	Store between 2-8°C. Use before expiry date on the label.		
Packing:	10/20/50 disposable plates.		

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal ortherapeutic use but for laboratory, diagnostic, research orfurther manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

Page 02 of 02

Rev: January2025