

TECHNICAL SHEET

PP034	Xylose Lysine Deoxycholate Agar Plate				
Formula					
Ingredients:		gms/lit.			
Yeast extract		3.00			
L-Lysine		5.00			
Lactose		7.50			
Sucrose		7.50			
Xylose		3.50			
Sodium chloride		5.00			
Sodium deoxycholate		2.50			
Sodium thiosulphate		6.80			
Ferric ammonium citrate		0.80			
Phenol red		0.08			
Agar		15.00			
Final pH (at 25°C): 7.4± 0.2					
Directions:					
Label the ready to use plate (PP034). Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.					
Principle:					
Xylose Lysine Deoxycholate Agar is a selective as well as differential medium. Yeast extract provides sources of nitrogen and carbon, as well as vitamins and cofactors required for growth. Xylose, lactose, and sucrose (Saccharose) are fermentable carbohydrates. Xylose is fermented by most enteric organisms except Shigella and Providencia. Lysine is added to differentiate Salmonella. As xylose is exhausted, Salmonella organisms decarboxylate lysine causing reversion to alkaline conditions. Alkaline reversion by other lysine – positive organisms is prevented by excess acid production from fermentation of lactose and sucrose. Sodium Thiosulfate and Ferric Ammonium citrate allow visualization of hydrogen sulfide production under alkaline conditions. Acidic conditions inhibit the reaction. Phenol red is pH indicator. Sodium chloride maintains osmotic balance in the medium. Agar is a solidifying agent. Sodium Deoxycholate in XLD agar inhibits growth of gram – positive organisms. This medium is an ideal medium for screening samples containing mixed flora of enteric pathogens as recovery of Salmonellae and Shigellae is not conspicuous by even profuse growth of other species.					
(I) QC Tests					
	pH:	7.4 ± 0.2			
	Color:	Red coloured medium.			
	Appearance:	Sterile Xylose Lysine Deoxycholate Agar in 85mm disposable plates.			
(II) Sterility test		Passes release criteria			
(III) Q.C. Test Microbiological					
Cultural characteristics observed after incubation at 35-37°C for 18-48 hours.					
	MICROORGANISM (ATCC)	INOCULUM (CFU)	GROWTH	RECOVERY	COLOR OF COLONY
	Salmonella typhimurium 14028	50 -100	LUXURIANT	≥50 %	red with black centers
	Salmonella Abony NCTC6017	50 -100	Good-luxuriant	≥50 %	red with black centers
	Escherichia coli 8739	50 -100	fair	20 -30 %	yellow
	Escherichia coli NCTC 9002	50 -100	fair	20 -30 %	yellow

TECHNICAL SHEET

Escherichia coli 25922	50 -100	fair	20 -30 %	yellow
Proteus vulgaris 13315	50 -100	Good-luxuriant	≥50 %	grey with black centers
Salmonella Paratyphi A9150	50 -100	Good-luxuriant	≥50 %	red
Salmonella Paratyphi B 8759	50 -100	Good-luxuriant	≥50 %	grey with black centers
Salmonella Enteritidis 13076	50 -100	Good-luxuriant	≥50 %	grey with black centers
Salmonella Typhi 6539	50 -100	Good-luxuriant	≥50 %	grey with black centers
Shigella flexneri12002	50 -100	fair-good	30 -40 %	red
Shigella dysenteriae13313	50 -100	Good-luxuriant	≥50 %	red
Shigella sonnei25931	50 -100	fair-good	30 -40 %	red
Enterobacter aerogenes13048	50 -100	fair	20 -30 %	yellow
Enterobacter cloacae 13047	50 -100	fair	20 -30 %	yellow
Staphylococcus aureus 6538	≥10 ³	inhibited	0%	-
Staphylococcus aureus25923	≥10 ³	inhibited	0%	-
Enterococcus faecalis 29212	≥10 ³	inhibited	0%	-

Refer disclaimer Overleaf

Precautions :	1. In Vitro diagnostic use only.
	2. Read the label before opening the container
Limitations :	1. Since the nutritional requirements of organisms vary, some strains may be encountered that fail to grow or grow poorly on this medium.
Use:	For the selection of Salmonella
Storage:	Store between 2-8°C. Use before expiry date on the label.
Packing:	20/50 disposable plates.

Disclaimer:

User must ensure suitability of the product(s) in their application prior to use. Products conform solely to the information contained in this and other related BIOMARKLABORATORIES publications.

The information contained in this publication is based on our in-house studies and market performance and is to the best of our knowledge true and accurate. BIOMARK LABORATORIES reserves the right to make changes to specifications and information related to the products at any time. Products are not intended for human or animal or therapeutic use but for laboratory, diagnostic, research or further manufacturing use only, unless otherwise specified. Statements contained herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.